

### Making Sequence logos

**Q1)** Below is a multiple alignment of 35 human sequences. The sequences have been aligned around a donor splice. That site is indicated as the boundary between the '**Dark blue**' and '**Dark red**' colours.

-----Exon|intron-----  
01234567890123456789  
**tatcacaATGGTAGGTAAC**  
TCAACCAGGAG**T**AAGTCTTG  
GTTGCACCC**T**GTAAAGTCTCA  
**tatcacaATGGTAGGTAAC**  
TCAACCAGGAG**T**AAGTCTTG  
CTTGCAGAG**G**TGTGACATG  
GCTCTACTCG**G**TAAGGTGAC  
GCCTGGAGAG**G**TAATGACCC  
CAAACCATT**G**TGAGTAATC  
GCCAGAGCAG**G**TAAAATATC  
GAACAGTCAG**G**TCTGTTGCT  
GAAGGCCAG**G**TGAGCATAA  
TCCTCTACAG**G**GTGGGTACAT  
GGCGTCCC**G**GTAAAGTATGG  
CCTCGTGCAG**G**TAAGATTAA  
TGCATGACAG**G**GTGAGTGTAA  
GAAATGTACAG**G**TAAGTCTCT  
GGTTCTCTGG**G**TAAGTAGAG  
AAATGTACAG**G**GTGAGTACTG  
ACCTCGCTT**G**GTACGTGGGA  
AATCAGACAG**G**GTATAGAAC  
AGGACAGAAG**G**TAATTTCT  
AACTATTGG**G**GTAGGTAGCA  
AAACTTGAAG**G**GTATGTTGTT  
CTGGGATAAG**G**TAAAAGTAT  
TTGCACCCAG**G**GTAGTGGAT  
ACTTCAATCG**G**GTATGTTTC  
ACAGAGAAA**G**TAACAACAA  
AATGGGAAAG**G**TAACAACAA  
CATGCTACAG**G**TAGGTGAAT  
ggctaggATGG**G**TGAGGGCGC  
CGACGCGGG**G**GTGAGAGGCG  
CATTGAGAAT**G**TGAGTTATT  
AACAGAGCAG**G**GTACTTGTAT  
TGAACCAAAG**G**TAAGACAT

Calculate the frequencies for positions **6–5**. You have each been assigned one column on the upper right corner of the handout.

position	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>	<b>0</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>
Counts A	15	12	20	4	0	0	23	23	5	6
Counts T	4	5	5	3	0	35	1	5	3	23
Counts C	8	13	4	2	0	0	1	2	1	1
Counts G	8	5	6	26	35	0	10	5	26	5
P(A)	0.43	0.34	0.57	0.11	0.0	0.0	0.66	0.66	0.14	0.17
P(T)	0.23	0.14	0.14	0.08	0.0	1.0	0.03	0.14	0.09	0.66
P(c)	0.11	0.37	0.11	0.05	0.0	0.0	0.03	0.06	0.03	0.03
P(G)	0.23	0.14	0.17	0.74	1.0	0.0	0.29	0.14	0.74	0.14

**Q2)** Calculate the Entropy and Information Content using the formula below

$$\text{Eq.1} \quad H(p) = -\sum_a p_a \log_2(p_a) = -\frac{1}{\log(2)} \sum_a p_a \log(p_a)$$

where  $\log_2$  is the logarithm with base 2, and  $\log$  is the logarithm with base 10 (or any base for that sake)

$$\text{Eq.2} \quad I = 2.0 - H(p)$$

position	6	7	8	9	0	1	2	3	4	5
Entropy	1.85	1.85	1.69	1.17	0	0	1.22	1.43	1.18	1.38
Information content	0.15	0.15	0.31	0.83	2	2	0.78	0.57	0.82	0.62

**Q3)** Where does the constant 2.0 come from in Eq.2?

**Q4)** Draw an approximate Logo Plot by hand on the White board

If you have internet-access

**Q5)** Submit the multiple alignment to the WebLogo server <http://weblogo.berkeley.edu/>

Make both the Logo plot and a frequency plot

Explain what you see on the two plots.